



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2020

Letter: Helicobacter pylori in proton pump inhibitor-associated biliary disease

Papaefthymiou, Apostolis ; Doulberis, Michael ; Polyzos, Stergios A ; Katsinelos, Panagiotis ; Liatsos, Christos ; Koffas, Apostolos ; Kazakos, Evangelos ; Deretzi, Georgia ; Srivastava, David S ; Kountouras, Jannis

DOI: <https://doi.org/10.1111/apt.15608>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-187557>

Journal Article

Accepted Version

Originally published at:

Papaefthymiou, Apostolis; Doulberis, Michael; Polyzos, Stergios A; Katsinelos, Panagiotis; Liatsos, Christos; Koffas, Apostolos; Kazakos, Evangelos; Deretzi, Georgia; Srivastava, David S; Kountouras, Jannis (2020). Letter: Helicobacter pylori in proton pump inhibitor-associated biliary disease. *Alimentary Pharmacology Therapeutics*, 51(2):313-314.

DOI: <https://doi.org/10.1111/apt.15608>

Letter: *Helicobacter pylori* in proton pump inhibitor-associated biliary disease

Apostolis Papaefthymiou¹ Michael Doulberis^{2,3} Stergios A. Polyzos^{3,4} Panagiotis Katsinelos³ Christos Liatsos¹ Apostolos Koffas⁵ Evangelos Kazakos³ Georgia Deretzi³, David S. Srivastava⁶ and Jannis Kountouras³

¹Department of Gastroenterology, 401 General Military Hospital of Athens, Athens, Greece

²Department of Gastroenterology and Hepatology, University of Zurich, Zurich, Switzerland

³Department of Internal Medicine, Second Medical Clinic, Ippokration Hospital, Aristotle University of Thessaloniki, Thessaloniki, Macedonia, Greece

⁴First Department of Pharmacology, School of Medicine, Aristotle University of Thessaloniki, Thessaloniki, Macedonia, Greece

⁵Department of Gastroenterology, Faculty of Medicine, School of Health Sciences, University of Thessaly, Larissa, Greece

⁶Department of Emergency Medicine, Inselspital, Bern University Hospital, University of Bern, Bern, Switzerland

Email: jannis@auth.gr; ancoratus2010@gmail.com

EDITORS,

Min et al,¹ in their cohort study, proposed the role of proton pump inhibitors (PPI) as a risk factor of cholangitis, especially during the period of their use. This observational result could include unrecognized confounders, one of which to consider being the overlap between *Helicobacter pylori* infection (*Hp-I*) and the conditions treated using a PPI; dyspepsia, peptic ulcer disease and, at least in some populations, gastro-oesophageal reflux disease—Barrett's oesophagus—oesophageal cancer sequence are associated with *Hp-I* and represent the most frequent indications for PPI prescription.² Moreover *Hp-I* is involved in PPI-related disorders discussed by the authors.¹⁻³

The impact of *Hp-I* with biliary tree diseases has been described. Bile stasis has primarily been connected with hepatobiliary *Hp* contamination, thus leading to a threefold risk of chronic lithiasis and cholecystitis, triggered by a perpetual inflammation of the gallbladder.^{4,5} In this respect, our data indicated *Hp* presence in gallbladder tissue in 19.33% of patients who were submitted to cholecystectomy due to lithiasis,⁶ while chronic biliary *Hp-I* has been identified in 75% of patients with gallbladder cancer.⁷

The proposed routes of contamination, although unproved, include ascending migration through the sphincter of Oddi and hematogenous spread to the liver and then excretion into bile. In this respect, the potential influx of activated monocytes infected with *Hp* (due to defective autophagy) from the circulation into the gallbladder might relate to gallbladder-related pathologies ("Trojan horse" pathway); biliary excretion of *Hp* invaded macrophages might trigger proinflammatory cytokine release, oxidative stress and inflammation, thus promoting gallstone formation and its

complications (cholecystitis/cholangitis/pancreatitis). Additionally, biliary *Hp-I* could provoke an imbalance of apoptotic cellular processes and contribute to biliary carcinogenesis. In vitro data support this, indicating that *Hp* promotes cell proliferation and suppresses apoptosis of human intrahepatic biliary epithelial cells by increasing reactive oxygen species release.⁸ Finally, the authors suggested as a possible pathogenetic mechanism of PPI-related cholangitis, the imbalance between the normal small intestine microbiota and the bacterial overgrowth due to gastric acid suppression.¹ Nevertheless, *Hp-I* seems to cause more significant alterations in gut microbiota than PPI use, thus contributing to biofilm formation.⁹

Hp-I has also been associated with a plethora of extragastric conditions, such as cardiovascular and neurodegenerative diseases, glaucoma, gastrointestinal tract oncogenesis, non-alcoholic fatty liver disease and hepatic encephalopathy.¹⁰ Therefore, it seems reasonable to consider adjusting for the presence of *Hp-I* as a potential cofounder in PPI-related disorders, as it is perhaps potentially relevant to a wide range of conditions necessitating their use.

ACKNOWLEDGEMENT

Declaration of personal interests: None. **FUNDING INFORMATION**

None.

LINKED CONTENT

This article is linked to Min et al paper. To view this article, visit <https://doi.org/10.1111/apt.15466>.

REFERENCES

1. Min YW, Kang D, Shin J-Y, et al. Use of proton pump inhibitors and the risk of cholangitis: a nationwide cohort study. *Aliment Pharmacol Ther.* 2019;50:760-768.
2. Kountouras J, Doulberis M, Papaefthymiou A, et al. A perspective on risk factors for esophageal adenocarcinoma: emphasis on *Helicobacter pylori* infection. *Ann N Y Acad Sci.* 2019;1452:12-17.
3. Kountouras J, Boziki M, Polyzos SA, et al. The emerging role of *Helicobacter pylori*-induced metabolic gastrointestinal dysmotility and neurodegeneration. *Curr Mol Med.* 2018;17:389-404.
4. Leong RWL, Sung JY. *Helicobacter* species and hepatobiliary diseases. *Aliment Pharmacol Ther.* 2002;16:1037-1045.
5. Cen L, Pan J, Zhou B, et al. *Helicobacter pylori* infection of the gall-bladder and the risk of chronic cholecystitis and cholelithiasis: a systematic review and meta-analysis. *Helicobacter.* 2018;23:e12457.
6. Kountouras J, Tsiaousi E, Trigonis S, et al. Comment on “The correlation of *Helicobacter pylori* with the development of cholelithiasis and cholecystitis: the results of a prospective clinical study in Saudi Arabia”. *Eur Rev Med Pharmacol Sci.* 2016;20:3-4.
7. Rawla P, Sunkara T, Thandra KC, Barsouk A. Epidemiology of gall-bladder cancer. *Clin Exp Hepatol.* 2019;5:93-102.
8. Ma F, Yang Y, Wang J-D, Quan Z-W, Zhou D. *Helicobacter pylori* and 17 β -estradiol induce human intrahepatic biliary epithelial cell abnormal proliferation and oxidative DNA damage. *Hepatobiliary Pancreat Dis Int.* 2017;16:519-527.
9. Parsons BN, Ijaz UZ, D’Amore R, et al. Comparison of the human gastric microbiota in hypochlorhydric states arising as a result of *Helicobacter pylori*-induced atrophic gastritis, autoimmune atrophic gastritis and proton pump inhibitor use. *PLOS Pathog.* 2017;13:e1006653.
10. Franceschi F, Gasbarrini A, Polyzos SA, Kountouras J. Extragastric diseases and *Helicobacter pylori*. *Helicobacter.* 2015;20:40-46.